

Analysis of Marijuana by Liquid Chromatographic Techniques A Literature Survey, 1990 - 2015

Robert F.X. Klein

U.S. Department of Justice
Drug Enforcement Administration
Special Testing and Research Laboratory
22624 Dulles Summit Court
Dulles, VA 20166
[robert.x.klein@usdoj.gov]

ABSTRACT: A survey of the liquid chromatographic analysis of marijuana, hemp, and related preparations (e.g., hashish, hash oil, “marijuana concentrates,” “cannabis smoke condensates,” etc.) for the major phytocannabinoids, as reported during the time frame 1990 through 2015,¹ is presented. 133 references are provided.

KEYWORDS: Marijuana, Cannabis, Hemp, Tetrahydrocannabinol, Phytocannabinoids, Liquid Chromatography, Forensic Chemistry.

Introduction

The qualitative and quantitative analysis of marijuana (*Cannabis sativa* L.) continues to be a significant task at most forensic laboratories. The most common techniques utilized for these analyses are GC/FID and GC/MS; however, while rapid and facile, these methodologies cannot handle thermally labile or non-volatile phytocannabinoids (such as Δ^9 -tetrahydrocannabinolic acid - A, THCA) without prior derivatization procedures - which tend to be time consuming, more expensive, and technique sensitive. An alternate approach for comprehensive analysis is to complement the GC-based methodology with a liquid chromatographic (LC-based) technique.

LC-based methodologies that have been utilized for this purpose range from basic (e.g., TLC with

UV or spray reagent detection) to highly sophisticated (e.g., UHPLC-MS/MS or UPC2). To the author’s knowledge, although this topic is lightly covered in several general reviews of the analysis of marijuana (e.g., 1,2,3), the last comprehensive survey of these methodologies appeared 30 years ago (4).

Search Details

Searches were conducted by the Chemical Abstracts Service’s Scientific & Technical Information Network (STN)[®], Google[®], PubMed, by reading select forensic journals, and/or by reviewing the reference citation lists of pertinent articles. The search terms for the analytical techniques included both their fully spelled-out names and their commonly utilized acronyms, as follows:

¹ Through November, 2015. Due to (normal) publication and/or abstracting delays, a small number of pre-2015 and 2015 dated references will not appear in this survey.

Automated Multiple Development (AMD);
 Capillary Electrophoresis (CE);
 Capillary Electrochromatography (CEC);
 Electrokinetic Chromatography (EKC);
 Hydrophilic Interaction Liquid Chromatography (HILIC);
 High Performance Liquid Chromatography (HPLC);
 High Performance Thin Layer Chromatography (HPTLC);
 Incremental Multiple Development (IMD);
 Liquid Chromatography (LC);
 Micellar Electrokinetic Capillary Chromatography (MECC);
 Optimal Performance Laminar Chromatography (OPLC, also known as overpressured layer chromatography or forced-flow TLC);
 Planar Chromatography (No Acronym);
 Reversed Phase High Performance Liquid Chromatography (RP-HPLC);
 Reversed Phase Liquid Chromatography (RP-LC);
 Supercritical Fluid Chromatography (SFC);
 Thin Layer Chromatography (TLC);
 Ultra High Performance Liquid Chromatography (UHPLC / UPLC);
 Ultra-Performance Convergence Chromatography (UPC2 / UPCC); and
 Two Dimensional Liquid Chromatography (2D-LC).

The search terms for marijuana included: Cannabinoids, cannabis, hash oil, hashish, hemp, hempseed, marihuana, marijuana, phyto-cannabinoids, tetrahydrocannabinol, and THC; street terms were not utilized. Analyses of rare / trace-level phytocannabinoids, or of phyto-cannabinoids and their metabolites in post-ingestion biological matrices, are not included in this survey. With the exception of a few reports that were re-published in *LCGC* or *American Laboratory*, “application notes,” printed “infomercials,” and similar publications also are not included. While there are no reasons to doubt

the validity of the presented analyses in these latter studies, virtually all such studies are either from scientific instrumentation companies touting the capabilities of one of their instruments or are from commercial (i.e., non-government) laboratories offering for-fee testing services for “medical” or “recreational” marijuana, marijuana concentrates, or “marijuana edibles,” and are not appropriate for this survey. Finally, references concerning the heterogeneity (inhomogeneity) of cannabis and the sampling protocols to address it - pertinent especially for the accurate quantitation of “herbal” cannabis and similar preparations - are covered as a separate category.

Results

References are organized first by general category (e.g., Planar Techniques), then by year (in reverse chronological order), and within year by author (in alphabetical order). Except for methodologies that are deemed (by this author) to be “advanced,” techniques that are interfaced with specialized detection methods are reported under the parent technique (e.g., HPLC-MS is reported under HPLC, but HPLC-MS/MS is reported under Advanced). References that report the use of more than one LC technique are reported under both/all categories. It should be noted that in many cases the LC technique is not the focus of the referenced article, but rather was used to complement or confirm the results acquired via a different methodology. Similarly, in many other cases the analysis of cannabis is only a minor aspect of a broader technique study.

Planar Techniques

2015 - by TLC/MS, for determination of cannabinoids and pesticides in cannabis (5); **2014** - by TLC, as a complement to a detailed study of the Duquenois-Levine color test (6); **2013** - by TLC (7); **2010** - by TLC, to analyze the components of ayurvedic (a folk remedy from the

Indian subcontinent that includes cannabis) (8); **2009** - by HPTLC, for the quantitation of THC and the qualitative analysis of other main neutral cannabinoids in cannabis (9); by TLC, as a comparison with and complement to a detailed study of *Salvia divinorum* and other salvia species (10); by TLC (from the UNODC Monograph on cannabis) (11); by TLC, for determination of cannabinoids in cannabis and hemp (12); **2007** - by TLC, for determination of cannabinoids in cannabis (13); by TLC (an effort to determine optimal mobile phases for select drugs, including cannabis) (14); **2006** - by TLC, as part of a more comprehensive analysis of cannabis (15); by TLC, for determination of cannabinoids in cannabis and hemp (16); **2005** - by TLC, as part of a more comprehensive analysis of cannabis (17); by TLC (from the SWGDRUG Monograph on cannabis) (18); by HPTLC, for determination of cannabinoids in both commercially available and cannabis oils stored long term (19); **2004** - by AMD, OPLC, and TLC (a comparative study of the analysis of cannabis by different planar chromatography techniques) (20); by TLC, for determination of cannabinoids in cannabis (21); **2003** - by OPLC (a technique study of a specialized OPLC instrument capable of simultaneously running 4 to 8 samples, including cannabis) (22); **2002** - by AMD, OPLC and TLC (a preliminary version of the work detailed in Reference #20, as conducted on different plants, including cannabis) (23); by TLC, for determination of cannabinoids in “monoecious” hemp (24); by OPLC, for determination of neutral cannabinoids in hemp (25); **1998** - by TLC, for analyses of cannabis resin and cannabis in unsmoked, handrolled cigarettes (26); by TLC, for the qualitative and quantitative analyses of cannabinoids in cannabis seeds (27); **1997** - by IMD (a technique study, including cannabis) (28); **1995** - by IMD (a feasibility study, including cannabis) (29); **1994** - by TLC, as part of a comprehensive chromatographic analysis of

cannabis (30); **1993** - by HPTLC (a technique study, including cannabis) (31); **1992** - by TLC, to determine the cannabinoid content of UK-grown plants (up to the 6th generation) from Moroccan, Sri Lankan, and Zambian seedstock (32); by HPTLC (a feasibility study, including cannabis) (33); by HPTLC, as part of the determination of cannabinoids in cannabis oil (34); **1990** - by TLC, for the analysis of derivatized cannabinoids (35); and by TLC, for the qualitative and quantitative analyses of cannabinoids in cannabis seeds (36).

Normal Phase LC / HPLC Techniques

2015 - by LC/MS, (37); by HPLC/DAD, for the analysis of cannabinoids and terpenes in cannabis (38); by HPLC, for the determination of 11 cannabinoids in biomass and in extracts of different varieties of cannabis (39); by HPLC and LC/MS (an overview of the marijuana testing rules in Colorado, the methods used for testing, and test results to date for “recreational marijuana”) (40); by HPLC (an overview of the marijuana testing rules in Colorado and the methods used for testing) (41); by HPLC/DAD (presenting two new, validated HPLC/DAD methods for identification and extract profiling based on the main patterns of cannabinoids and other phenolics in cannabis) (42); by HPLC, for the determination of the relative percentage of THCA and THC in cannabis, and the impact of different storage temperatures on stability (43); by LC/MS, as a complement to a DNA genotyping study (44); by HPLC, for determination of cannabinoids in “marijuana edibles” (45); by HPLC, for determination of cannabinoids in “marijuana edibles” (a feasibility study with spiked samples) (46); **2014** - by HPLC, for identification and quantitation of cannabinoids in cannabis (47); by HPLC/DAD, for quantitation of THC, THCA, CBN, and CBD in seized cannabis (48); by HPLC, to determine THC and THCA (49); by HPLC/DAD, to determine THC in hempseed oil (50); **2013** - by HPLC, for

determination of THC following cloud point extraction of cannabis resin (51); by LC/MS, for analysis of cannabinoids in laser-microdissected trichomes of “medicinal” cannabis (52); by HPLC, for determination of potency and cannabinoid profiles (53); **2012** - by HPLC, as a complement to the voltametric determination of THC (54); by HPLC/DAD, for the determination of THC and other major cannabinoids in cannabis cuttings and seedlings during plant growth (55); by HPLC with chemiluminescence detection, for determination of CBD in industrial-grade hemp (stated to be applicable for determination of THC in cannabis) (56); by Nano-LC, for determination of synthetic cannabinoids and THC in herbal blends (focus is on synthetic cannabinoids) (57); by LC/MS (a review of the use of LC/MS for the detection and quantitation of cannabinoids) (58); by HPLC, for the determination of cannabinoids in cannabis oil during long-term storage (59); by HPLC, for the determination of cannabinoids in cannabis resin during long-term storage (60); by HPLC, for the identification and characterization of “special types of herbal cannabis” (61); by HPLC (a study to determine the optimal solvent and conditions for extraction of THC, CBD, and CBN from cannabis resin) (62); by HPLC, for determination of THC in cannabis (63); **2011** - by HPLC, for the determination of cannabinoids in cannabis during different storage conditions (64); by HPLC/DAD, for determination of THC and (separately) THCA from cannabis after isolation via two different flash chromatography systems (65); **2010** - by HPLC/DAD, for monitoring the long term stability of cannabis resin and cannabis extracts (66); **2009** - by HPLC/DAD, for the qualitative and quantitative determination of the major cannabinoids in cannabis (67); by HPLC, as a complement to an HPTLC study (68); by LC/MS, as a complement to a DART-TOF-MS-based screening for THC in cannabis (69); by High-Temperature LC (a technique study, including analysis of THC in cannabis) (70); by

HPLC, to determine cannabinoids in vaporized cannabis (71); by HPLC (from the UNODC Monograph on cannabis) (72); by HPLC, to determine the effects of tobacco on the levels of cannabinoids in vaporized cannabis (73); **2008** - by HPLC, to determine the effects of different preparation methods on the levels of cannabinoids in vaporized cannabis (74); **2007** - by HPLC, for the determination of the major cannabinoids in “medicinal grade” cannabis (75); by HPLC, for the determination of the major cannabinoids in cannabis tea (76); by HPLC, for the determination of cannabinoids in hemp (77); **2006** - by HPLC, for the determination of the major cannabinoids in “medicinal grade” cannabis (78); by HPLC, as a complement to a DNA study of drug-type versus fiber-type cannabis (79); **2005** - by HPLC/DAD and HPLC with fluorescence detection (as part of a detailed chromatographic and spectroscopic analysis of the cannabinoids in cannabis) (80); by HPLC at elevated pressure (a technique study, including analysis of THC in cannabis) (81); by HPLC/DAD, for analysis of the cannabinoids in cannabis (82); **2004** - by HPLC, for analysis of the cannabinoids in cannabis (83); by HPLC/ESI-MS (a technique study, including analysis of the cannabinoids in cannabis) (84); **2003** - by HPLC, for determination of CBD in hempseed oil (85); **2002** - HPLC, for the determination of cannabinoids in cannabis (86); by HPLC, as a complement to a DNA (ISSR) study (87); by HPLC (an optimization study of the HPLC separation conditions for cannabinoids in cannabis, including Δ^9 - versus Δ^8 -THC) (88); **2001** - by HPLC, as a complement to a study of the supercritical fluid extraction of cannabis (89); **2000** - by Capillary LC with electrochemical detection, for determination of the cannabinoids in cannabis (90); by HPLC, for determination of the cannabinoids in hashish (91); by HPLC, for determination of the cannabinoids in cannabis (92); by HPLC with UV or fluorescence detection, for determination of THC and THCA in hemp-

containing foods (93); **1998** - by HPLC, for analysis of cannabinoids in hemp (94); **1997** - by HPLC, for the qualitative and quantitative analyses of cannabinoids in cannabis (95); **1996** - by LC/MS, for determination of the cannabinoids in hashish (96); **1995** - by HPLC, as a complement to a DNA (RAPD) study of different samples of cannabis (97); by HPLC/DAD, for the qualitative and quantitative determination of neutral and acidic cannabinoids in cannabis (for profiling purposes) (98); **1994** - by HPLC, as part of a comprehensive chromatographic analysis of cannabis (99); by HPLC, as a complement to a study of the supercritical fluid extraction of cannabis and hashish (100); **1992** - by HPLC, to determine the cannabinoid content of UK-grown plants (up to the 6th generation) from Moroccan, Sri Lankan, and Zambian seedstock (101); **1991** - by HPLC, to determine the uniformity of hashish samples (102); by HPLC (as a complement to an MECC technique study, including analysis of cannabis) (103); and **1990** - by HPLC with UV or fluorescence detection, following derivatization (a technique study, including analysis of cannabis) (104).

Reversed Phase LC and HPLC Techniques

2015 - by RP-HPLC (on three different columns), for determination of the cannabinoids in cannabis and a marijuana concentrate (105); **1996** - by RP-HPLC, for analysis of “drugs of abuse,” including cannabis (106); **1994** - by RP-HPLC (a technique study, including analysis of THC in cannabis) (107); **1993** - by RP-HPLC, for determination of THC in cannabis (108); and **1990** - by RP-HPLC, to characterize the lipophilicity of natural and synthetic analogs of THC (109).

Electrokinetic Techniques

2004 - by CE/ESI-MS (a technique study, also covering HPLC/ESI-MS, including analysis of the

cannabinoids in cannabis) (110); **1998** - by CEC (a technique study, including analysis of THC in cannabis) (111); by CEC/DAD, for the determination of the cannabinoids in cannabis (112); and **1991** - by MECC, for the determination of the cannabinoids in cannabis (113).

Advanced Techniques

2015 - by HILIC, as part of an RP-HPLC study for the determination of the cannabinoids in cannabis and a marijuana concentrate (114); by 2D-LC with chemiluminescence detection, for screening of cannabinoids in industrial-grade hemp (115); by EI-LC/MS with supersonic molecular beams (an introductory technique study, including cannabis) (116); by UPC2, for the determination of the cannabinoids in cannabis (117); **2014** - by HPLC/MS and HPLC-MS/MS, for identification and quantitation of cannabinoids in cannabis (118); **2013** - by UHPLC/MS (a technique review, including analysis of a mixture of drug standards including THC, CBD, and CBN, plus analyses of two different baked goods that contained THC) (119); by LC-MS/MS, for screening of “botanicals” (including cannabis) in food supplements (120); by UHPLC-MS/MS, for determination of THC in hemp food (121); by UHPLC-Q-ToF-MS/MS, for determination of cannabinoids in hemp seed pills (a traditional Chinese medication) (122); **2012** - by LC/ESI-MS/MS, to investigate the isomerization of CBD and THC during positive ESI analyses (123); **2011** - by HILIC (a technique study, including analysis of THC in cannabis) (124); by LC-MS/MS, for determination of cannabinoids in industrial hemp (125); by UPLC-MS/MS, for determination of THC in edible vegetable oil (126); **2010** - by UHPLC/MS (a technique study, including analysis of THC in cannabis) (127); by UPLC-MS/MS, for determination of cannabinoids in edible oil (128); **2009** - by LC/MS and LC-MS/MS, to evaluate microwave-assisted derivatization of THC (129); by UHPLC/MS, for

determination of cannabinoids in baked goods (130); **2004** - by LC-Ion Trap-MS/MS, for analysis of the cannabinoids in cannabis (131); **2000** - by SFC, for the determination of the cannabinoids in cannabis (132); by SFC (a technique study, including analysis of THC in cannabis) (133); **1998** - by LC with thermospray-MS detection and by LC-MS/MS, for determination of the cannabinoids in cannabis (134); **1997** - by SFC with APCI-MS detection, for analysis of THC in cannabis (135); and **1993** - by HPLC with post-elution photoirradiation followed by DAD or thermospray MS detection (a technique study, including analysis of THC in cannabis) (136).

Heterogeneity (Inhomogeneity) of Cannabis and Sampling Protocols to Reduce its Impact

Although the “within sample” heterogeneity of “herbal” cannabis and the resulting, inherent variability on their quantitative analyses are widely recognized (e.g., 137,138), to date there

have been few studies that specifically addressed these issues (139,140). Reflecting the importance of this topic, however, a comprehensive, three-part study was recently published (141,142,143). The heterogeneity of all plant materials utilized for medicinal, nutritional, or similar purposes has also been addressed in more general terms by the U.S. Food and Drug Administration (144) and the World Health Organization (145,146).

Acknowledgment

The assistance of DEA Librarian Kristin Carr in acquiring numerous references is gratefully acknowledged.

References

[Note: In order to avoid the occasional overly wide spacings created by the use of fully justified columns, the references are provided in full page, left-justified format.]

1. Thomas BF, ElSohly MA. The Analytical Chemistry of Cannabis: Quality Assessment, Assurance, and Regulation of Medicinal Marijuana and Cannabinoid Preparations. Elsevier, Waltham, Massachusetts:2016. [Note: Appeared on-line in late 2015.]
2. Brenneisen R. Chemistry and Analysis of Phytocannabinoids and other Cannabis Constituents. In: Marijuana and the Cannabinoids. ElSohly MA, Editor. Humana Press Inc., Totowa, New Jersey:2007.
3. Raharjo TJ, Verpoorte R. Methods for the analysis of cannabinoids in biological materials: A review. *Phytochemical Analysis* 2004;15(2):79-94.
4. Vollner L, Bieniek D, Korte F. Review of analytical methods for identification and quantification of cannabis products. *Regulatory Toxicology and Pharmacology* 1986;6(4):348-358.
5. Eikel D, Prosser SJ, Henion JD. Using compact mass spectrometry for detection and quantification of cannabis-related compounds. *LCGC Europe* 2015;(Suppl.):20-27.
6. Cantrell S, McGill J. Duquenois-Levine test revisited: Studies toward reviewable data. Abstracts, 66th Southeast Regional Meeting of the American Chemical Society, Nashville, TN, United States, October 16-19, 2014; SERMACS-798.

7. Srivastava A, Yadav VK. Microscopical and chemical study of *Cannabis sativa*. Journal of Forensic Research 2013;5(1):210 (6 pages).
8. Mohammad A, Sharma S, Bhawani SA, Singh RB. Identification and separation of *Cannabis sativa*, *Embllica ribes*, *Myristica fragrans* and *Piper longum* from organic extract on silica gel surface with anionic micellar solvent system: Application in ayurvedic medicine. Open Nutraceuticals Journal 2010;3:112-118.
9. Fishedick JT, Glas R, Hazekamp A, Verpoorte R. A qualitative and quantitative HPTLC densitometry method for the analysis of cannabinoids in *Cannabis sativa* L. Phytochemical Analysis 2009;20(5):421-426.
10. Jermain JD, Evans HK. Analyzing *Salvia divinorum* and its active ingredient salvinorin A utilizing thin layer chromatography and gas chromatography/mass spectrometry. Journal of Forensic Sciences 2009;54(3):612-616.
11. UNODC. Recommended methods for the identification and analysis of cannabis and cannabis products. United Nations Office on Drugs and Crime: Vienna, 2009. [Note: TLC in Section 5.4.5, pps. 36-38.]
12. Zelenina ON, Serkov VA. Methods for assessment of cannabinoid contents in hemp plants. Vestnik Rossiiskoi Akademii Sel'skokhozyaistvennykh Nauk 2009;(2):53-55.
13. Bladek J, Polak P, Jarzyna E, Jakubowska I. Analysis of opiates and cannabinols by TLC. Biuletyn Wojskowej Akademii Technicznej 2007;56(3):257-267.
14. Kuila DK, Lahiri SC. Search for suitable mobile phase in TLC analysis of different drugs of forensic interest and their gas liquid chromatographic experiment. Journal of the Indian Chemical Society 2007;84(1):69-73.
15. Wachowiak R, Strach B. Analysis of active components of evidence materials secured in the cases of drugs abuse associated with amphetamines and cannabis. Archiwum Medycyny Sadowej i Kryminologii 2006;56(4):251-257. [Note: TLC is only a minor aspect of the presented analyses.]
16. Zelenina ON. Assessment of cannabinoid content in hemp plants. Vestnik Rossiiskoi Akademii Sel'skokhozyaistvennykh Nauk 2006;(6):43-45.
17. Hazekamp A, Peltenburg A, Verpoorte R, Giroud C. Chromatographic and spectroscopic data of cannabinoids from *Cannabis sativa* L. Journal of Liquid Chromatography & Related Technologies 2005;28(15):2361-2382.
18. SWGDRUG. In: Monograph on Marijuana; Section 3.3, Thin Layer Chromatography. May 13, 2005.
19. Yotoriyama M, Ishiharajima E, Kato Y, Nagato A, Sekita S, Watanabe K, Yamamoto I. Identification and determination of cannabinoids in both commercially available and cannabis oils stored long term. Journal of Health Science 2005;51(4):483-487.

20. Galand N, Ernouf D, Montigny F, Dollet J, Pothier J. Separation and identification of cannabis components by different planar chromatography techniques (TLC, AMD, OPLC). *Journal of Chromatographic Science* 2004;42(3):130-134.
21. See Raharjo and Verpoorte, Reference #3.
22. Papillard D, Manach M, Laroche S, Truchy C, Bryson N, Mincsovcics E. Simultaneous, parallel purification of complex samples on an unsegmented flat column using a 4-channel multi-OPLC. *LCCG Europe* 2003;16(12a):60-61.
23. Galand N, Pothier J, Viel C. Plant drug analysis by planar chromatography. *Journal of Chromatographic Science* 2002;40(10):585-597.
24. Laiko IM. The methods of cannabinoid content evaluation in monoecious hemp. *Sel'skokhozyaistvennaya Biologiya* 2002;(3):117-120.
25. Szabady B, Hidvegi E, Nyiredy S. Determination of neutral cannabinoids in hemp samples by overpressured-layer chromatography. *Chromatographia* 2002;56(Suppl.):S165-S168.
26. Buchanan BE, O'Connell D. Survey on cannabis resin and cannabis in unsmoked handrolled cigarettes seized in the Republic of Ireland. *Science & Justice* 1998;38(4):221-224.
27. Matsunaga T. Discrimination of cannabis seeds. *Japanese Journal of Forensic Toxicology* 1998;16(2):124-127.
28. Szabady B, Ruzinko M, Nyiredy S. Prediction of retention data when using incremental multiple development techniques. *Chromatographia* 1997;45:369-372.
29. Szabady B, Ruzinkó M, Nyiredy. Prediction of retention data in multiple development. Part 1. Linearly increasing development distances. *JPC - Journal of Planar Chromatography - Modern TLC* 1995;8(4):279-283.
30. Debruyne D, Albessard F, Bigot MC, Moulin M. Comparison of three advanced chromatographic techniques for cannabis identification. *Bulletin on Narcotics* 1994;46(2):109-121.
31. Pothier J, Galand N, Viel C. Determination of some narcotic and toxic alkaloidal compounds by overpressured thin-layer chromatography with ethyl acetate as eluent. *Journal of Chromatography* 1993;634(2):356-359.
32. Pitts JE, Neal JD, Gough TA. Some features of cannabis plants grown in the United Kingdom from seeds of known origin. *Journal of Pharmacy and Pharmacology* 1992;44(12):947-951.
33. Pothier J, Galand N, Viel C. Rapid characterization of narcotic and toxic compounds by high-pressure thin-layer chromatography. *Journal de Toxicologie Clinique et Experimentale* 1992;12(8):495-501.
34. Yotoriyama M, Okada Y, Suzuki K. Determination of cannabinoids in cannabis oil by capillary gas chromatography. *Journal of Toxicology and Environmental Health* 1992;38(5):471-475.

35. Lavanya K, Baggi TR. An improved thin-layer chromatographic method for the detection and identification of cannabinoids in cannabis. *Forensic Science International* 1990;47(2):165-71.
36. Matsunaga T, Nagatomo H, Yamamoto I, Yoshimura H. Qualitative and quantitative analyses of cannabinoids in cannabis seeds. *Japanese Journal of Forensic Toxicology* 1990;8(2):88-89.
37. See Eikel, Prosser, *et al.*, Reference #5.
38. Giese MW, Lewis MA, Giese L, Smith KM. Development and validation of a reliable and robust method for the analysis of cannabinoids and terpenes in cannabis. *Journal of the AOAC International* 2015;98(6):1503-1522.
39. Gul W, Gul SW, Radwan MM, Wanas AS, Mehmedic Z, Khan II, Sharaf MH, ElSohly MA. Determination of 11 cannabinoids in biomass and extracts of different varieties of cannabis using high-performance liquid chromatography. *Journal of the AOAC International* 2015;98(6):1523-1528.
40. LaFrate A. State mandated testing of retail marijuana in Colorado. Abstracts of Papers, 249th ACS National Meeting & Exposition, Denver, CO, United States, March 22-26, 2015; CHAS-4.
41. LaFrate A. Potency and contaminant testing of retail marijuana in Colorado. Abstracts of Papers, 249th ACS National Meeting & Exposition, Denver, CO, United States, March 22-26, 2015; CHAS-9.
42. Peschel W, Politi M. ¹H NMR and HPLC/DAD for *Cannabis sativa* L. chemotype distinction, extract profiling and specification. *Talanta* 2015;140:150-65.
43. Taschwer M, Schmid MG. Determination of the relative percentage distribution of THCA and Δ(9)-THC in herbal cannabis seized in Austria - Impact of different storage temperatures on stability. *Forensic Science International* 2015;254:167-171.
44. Welling MT, Liu L, Shapter T, Raymond CA, King GJ. Characterisation of cannabinoid composition in a diverse *Cannabis sativa* L. germplasm collection. *Euphytica* 2015: Ahead of Print.
45. Vandrey R, Raber JC, Raber ME, Douglass B, Miller C, Bonn-Miller MO. Cannabinoid dose and label accuracy in edible medical cannabis products. *Journal of the American Medical Association* 2015;313(24):2491-2493.
46. Wang X, Fanning K. Determination of 35 pesticides and 3 cannabinoids in marijuana edibles. LCGC North America, Nov 20, 2015. [Note: Also appears in the LCGC Europe issue of the same date.]
47. Ahmada UK, Muniandy Y, Hassan MS. Physical analysis and chemical profiling of illicit herbal cannabis using multivariate analysis. *Malaysian Journal of Forensic Sciences* 2014;5(1):26-34.
48. Ambach L, Penitschka F, Broillet A, König S, Weinmann W, Bernhard W. Simultaneous quantification of delta-9-THC, THC-acid A, CBN and CBD in seized drugs using HPLC-DAD. *Forensic Science International* 2014;243C:107-111.

49. Rymanowski M. Cannabis - Review of the issues related to determination of the total content of delta-9-tetrahydrocannabinol (Δ -9-THC) and delta 9-tetrahydrocannabinolic acid (Δ -9-THCA-A). *Problemy Kryminalistyki* 2014;285(3):1-22.
50. Shishovska M, Doneva D, Starkovska K, Arsova-Sarafinovska Z. Determination of Δ -9-tetrahydrocannabinol by HPLC/DAD in food supplement samples of hempseed oil. Presented at: XXIII Congress of Chemists and Technologists of Macedonia, Ohrid, Macedonia, October 8-11, 2014; AC-018.
51. Ameer S, Haddou B, Derriche Z, Canselier JP, Gourdon C. Cloud point extraction of Δ -9-tetrahydrocannabinol from cannabis resin. *Analytical and Bioanalytical Chemistry* 2013;405(10):3117-3123.
52. Happyana N, Agnolet S, Muntendam R, Van Dam A, Schneider B, Kayser O. Analysis of cannabinoids in laser-microdissected trichomes of medicinal *Cannabis sativa* using LCMS and cryogenic NMR. *Phytochemistry* 2013;87:51-59.
53. Swift W, Wong A, Li KM, Arnold JC, McGregor IS. Analysis of cannabis seizures in NSW, Australia: Cannabis potency and cannabinoid profile. *PLoS One* 2013;8(7):e70052.
54. Balbino MA, de Menezes MM, Eleotério IC, Saczk AA, Okumura LL, Tristão HM, de Oliveira MF. Voltammetric determination of Δ -9-THC in glassy carbon electrode: An important contribution to forensic electroanalysis. *Forensic Science International* 2012;221(1-3):29-32.
55. De Backer B, Maebe K, Verstraete AG, Charlier C. Evolution of the content of THC and other major cannabinoids in drug-type cannabis cuttings and seedlings during growth of plants. *Journal of Forensic Sciences* 2012;57(4):918-922.
56. Holland BJ, Francis PS, Li B, Tsuzuk Ti, Adcock JL, Barnett NW, Conlan XA. Chemiluminescence detection of cannabinoids and related compounds with acidic potassium permanganate. *Drug Testing and Analysis* 2012;4(7-8):675-679.
57. Merola G, Aturki Z, D'Orazio G, Gottardo R, Macchia T, Tagliaro F, Fanali S. Analysis of synthetic cannabinoids in herbal blends by means of nano-liquid chromatography. *Journal of Pharmaceutical and Biomedical Analysis* 2012;71:45-53.
58. Quintela O, Crouch DJ. The determination of cannabinoids using liquid chromatography with mass spectrometric detection. *Methods in Molecular Biology* 2012;902(LC-MS in Drug Analysis):75-90.
59. Trofin IG, Dabija G, Vaireanu D-I, Filipescu L. Long-term storage and cannabis oil stability. *Revista de Chimie* 2012;63(3):293-297.
60. Trofin IG, Dabija G, Vaireanu D-I, Filipescu L. The influence of long-term storage conditions on the stability of cannabinoids derived from cannabis resin. *Revista de Chimie* 2012;63(4):422-427.
61. Trofin IG, Vlad CC, Noja VV, Dabija G. Identification and characterization of special types of herbal cannabis. *Scientific Bulletin - University "Politehnica" of Bucharest, Series B: Chemistry and Materials Science* 2012;74(1):119-130.

62. Zhai W-f. Study on pre-treatment methods of cannabis resin in the public security cases. *Huaxue Gongchengshi* 2012;26(8):7-9.
63. Zhai W-f, Zhang C-s, Gao L-s. Study on determination of Δ^9 -tetrahydrocannabinol in cannabis by high performance liquid chromatography. *Fenxi Ceshi Xuebao* 2012;31(11):1379-1384.
64. Trofin IG, Vlad CC, Dabija G, Filipescu L. Influence of storage conditions on the chemical potency of herbal cannabis. *Revista de Chimie* 2011;62(6):639-645.
65. Wohlfarth A, Mahler H, Auwaerter V. Rapid isolation procedure for Δ^9 -tetrahydrocannabinolic acid A (THCA) from *Cannabis sativa* using two flash chromatography systems. *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences* 2011;879(28):3059-3064.
66. Lindholst C. Long term stability of cannabis resin and cannabis extracts. *Australian Journal of Forensic Sciences* 2010;42(3):181-190.
67. De Backer B, Debrus B, Lebrun P, Theunis L, Dubois N, Decock L, Verstraete A, Hubert P, Charlier C. Innovative development and validation of an HPLC/DAD method for the qualitative and quantitative determination of major cannabinoids in cannabis plant material. *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences* 2009;877(32):4115-4124.
68. See Fishedick, Glas, *et al.*, Reference #9.
69. Kawamura M, Kikura-Hanajiri R, Goda Y. Simple and rapid screening for psychotropic natural products using direct analysis in real time (DART)-TOFMS. *Yakugaku Zasshi* 2009;129(6):719-725.
70. Lurie IS, Li L. Use of High-Temperature Liquid Chromatography with sub-2 μ m particle C18 columns for the analysis of seized drugs. *Journal of Liquid Chromatography & Related Technologies* 2009;32(18):2615-2626.
71. Pomahacova B, Van der Kooy F, Verpoorte R. Cannabis smoke condensate III: The cannabinoid content of vaporised *Cannabis sativa*. *Inhalation Toxicology* 2009;21(13):1108-1112.
72. See UNODC, Reference #11. [Note: HPLC in Section 5.4.8, pps. 41-43.]
73. Van der Kooy F, Pomahacova B, Verpoorte R. Cannabis smoke condensate II: Influence of tobacco on tetrahydrocannabinol levels. *Inhalation Toxicology* 2009;21(2):87-90.
74. Van der Kooy F, Pomahacova B, Verpoorte R. Cannabis smoke condensate I: The effect of different preparation methods on tetrahydrocannabinol levels. *Inhalation Toxicology* 2008;20(9):801-804.
75. Hazekamp A. Development and validation of an HPLC method for the determination of major cannabinoids from medicinal grade *Cannabis sativa* plant material. Chapter 7 (pps. 91-105) in *Cannabis - Extracting the medicine*. Doctoral Thesis, Department of Pharmacognosy, Leiden University, The Netherlands, September 5, 2007.]

76. Hazekamp A, Bastola K, Rashidi H, Bender J, Verpoorte R. Cannabis tea revisited: A systematic evaluation of the cannabinoid composition of cannabis tea. *Journal of Ethnopharmacology* 2007;113(1):85-90. [Note: This article is reprinted as Chapter 8 (pps. 107-117) in *Cannabis - Extracting the medicine*. Doctoral Thesis, Department of Pharmacognosy, Leiden University, The Netherlands, September 5, 2007.]
77. Stanaszek R, Zuba D. A comparison of developed and validated chromatographic methods (HPLC, GC-MS) for determination of delta-9-tetrahydrocannabinol (delta-9-THC) and delta-9-tetrahydrocannabinolic acid (delta-9-THCA-A) in hemp. *Z Zagadnien Nauk Sadowych* 2007;71:313-322.
78. Hazekamp A. An evaluation of the quality of medicinal grade cannabis in the Netherlands. *Cannabinoids* 2006;1(1):1-9.
79. Kojoma M, Seki H, Yoshida S, Muranaka T. DNA polymorphisms in the tetrahydrocannabinolic acid (THCA) synthase gene in "drug-type" and "fiber-type" *Cannabis sativa* L. *Forensic Science International* 2006;159(2-3):132-140.
80. See Hazekamp, Peltenburg, *et al.*, Reference #17.
81. Lurie IS. High-performance liquid chromatography of seized drugs at elevated pressure with 1.7 μ m hybrid C18 stationary phase columns. *Journal of Chromatography A* 2005;1100(2):168-175.
82. Stambouli H, El Bouri A, Bellimam MA, Bouayoun T, El Karni N. Cultivation of *Cannabis sativa* L. in northern Morocco. *Bulletin on Narcotics* 2005;57(1-2):79-118.
83. See Raharjo and Verpoorte, Reference #3.
84. Smyth WF, Brooks P. A critical evaluation of high performance liquid chromatography-electrospray ionisation-mass spectrometry and capillary electrophoresis-electrospray-mass spectrometry for the detection and determination of small molecules of significance in clinical and forensic science. *Electrophoresis* 2004;25(10-11):1413-1446.
85. Zhang G, Guo J, Bi K. Determination of cannabidiol in hempseed oil by HPLC. *Zhongcaoyao* 2003;34(5):415-417.
86. Gambaro V, Dell'Acqua L, Fare F, Frolidi R, Saligari E, Tassoni G. Determination of primary active constituents in cannabis preparations by high-resolution gas chromatography/flame ionization detection and high-performance liquid chromatography/UV detection. *Analytica Chimica Acta* 2002;468(2):245-254.
87. Kojoma M, Iida O, Makino Y, Sekita S, Satake M. DNA fingerprinting of *Cannabis sativa* using inter-simple sequence repeat (ISSR) amplification. *Planta Medica* 2002;68(1):60-63.
88. Wojtasik E, Anyzewska M, Arent I. The optimization of the separation conditions for cannabinoids from *Cannabis sativa* L. var. Indica and application of the method to determine the content of delta(9)-tetrahydrocannabinol in plant material. *Journal of Liquid Chromatography and Related Technologies* 2002;25(6):949-959.

89. Eory L, Danos B, Veress T. Supercritical fluid extraction of tetrahydrocannabinol from marijuana study of the effect of particle size. *Z Zagadnien Nauk Sadowych* 2001;47:322-327.
90. Backofen U, Hoffmann W, Matysik FM. Determination of cannabinoids by capillary liquid chromatography with electrochemical detection. *Biomedical Chromatography* 2000;14(1):49-52.
91. Ferioli V, Rustichelli C, Pavesi G, Gamberini G. Analytical characterisation of hashish samples. *Chromatographia* 2000;52(1):39-44.
92. Tsatsakis AM, Tutudaki M, Stiakakis I, Dimopoulou M, Tzatzarakis M, Michalodimitrakis M. Characterization of cannabis plants phenotypes from illegal cultivations in Crete. *Bollettino Chimico Farmaceutico* 2000;139(3):140-145.
93. Zoller O, Rhyn P, Zimmerli B. High-performance liquid chromatographic determination of delta-9-tetrahydrocannabinol and the corresponding acid in hemp containing foods with special regard to the fluorescence properties of delta-9-tetrahydrocannabinol. *Journal of Chromatography A* 2000;872(1-2):101-110.
94. Rustichelli C, Ferioli V, Baraldi M, Zanolì P, Gamberini G. Analysis of cannabinoids in fiber hemp plant varieties (*Cannabis sativa*) by high-performance liquid chromatography. *Chromatographia* 1998;48(3/4):215-222.
95. Chang SG, Chen CY, Chin FS, Chang HC, Li JH. Qualitative and quantitative analyses of cannabinoids in *Cannabis sativa*, huoma-jen and *Agave sisalana*. *Chinese Pharmaceutical Journal* 1997;49(2):77-88.
96. Rustichelli C, Ferioli V, Vezzalini F, Rossi MC, Gamberini G. Simultaneous separation and identification of hashish constituents by coupled liquid chromatography-mass spectrometry (HPLC-MS). *Chromatographia* 1996;43(3/4):129-134.
97. Gillan R, Cole MD, Linacre A, Thorpe JW, Watson ND. Comparison of *Cannabis sativa* by random amplification of polymorphic DNA (RAPD) and HPLC of cannabinoids: A preliminary study. *Science & Justice* 1995;35(3):169-177.
98. Lehmann T, Brenneisen R. High performance liquid chromatographic profiling of cannabis products. *Journal of Liquid Chromatography* 1995;18(4):689-700.
99. See Debruyne, Albessard, *et al.*, Reference #30.
100. Veress T. Sample preparation by supercritical fluid extraction for quantification. A model based on the diffusion-layer theory for determination of extraction time. *Journal of Chromatography A* 1994;668(2):285-291.
101. See Pitts, Neal, *et al.*, Reference # 32.
102. Kovar K-A, Linder H. Investigation of hashish: Content uniformity of different samples by coupled HPLC/PC-analysis. *Archiv der Pharmazie* 1991;324(6):329-333.

103. Weinberger R, Lurie IS. Micellar electrokinetic capillary chromatography of illicit drug substances. *Analytical Chemistry* 1991;63(8):823-837.
104. Moore JM. The application of chemical derivatization in forensic drug chemistry for gas and high-performance liquid chromatographic methods of analysis. *Forensic Science Review* 1990;2(2):79-124.
105. Hung C-H, Zukowski J, Jensen DS, Miles AJ, Sulak C, Dadson AE, Linford MR. Separation of cannabinoids on three different mixed-mode columns containing carbon/nanodiamond/amine-polymer superficially porous particles. *Journal of Separation Science* 2015;38(17):2968-2974.
106. Ma C, Duan HJ, Xu YQ, Zhou TH. Reversed-phase high-performance liquid chromatographic determination of drugs of abuse. *Chinese Chemical Letters* 1996;7(11):1027-1028.
107. Bogusz M, Erkens M. Reversed-phase high-performance liquid chromatographic database of retention indices and UV spectra of toxicologically relevant substances and its interlaboratory use. *Journal of Chromatography A* 1994;674(1-2):97-126.
108. Ghysel MH, Salvadore O, Berquez N, Deramaux C, Vilette JM, Haguenoer JM. Chromatographic determination of Δ -9-tetrahydrocannabinol in new forms of cannabis. *Analisis* 1993;21(6):M31-M33.
109. Thomas BF, Compton DR, Martin BR. Characterization of the lipophilicity of natural and synthetic analogs of Δ 9-tetrahydrocannabinol and its relationship to pharmacological potency. *Journal of Pharmacology and Experimental Therapeutics* 1990;255(2):624-630.
110. See Smyth and Brooks, Reference #84.
111. Lurie IS, Conver TS, Ford VL. Simultaneous separation of acidic, basic, and neutral organic compounds, including strong and moderate acids and bases, by capillary electrochromatography. *Analytical Chemistry* 1998;70(21):4563-4569.
112. Lurie IS, Meyers RP, Conver TS. Capillary electrochromatography of cannabinoids. *Analytical Chemistry* 1998;70(15):3255-3260.
113. See Weinberger and Lurie, Reference #103.
114. See Hung, Zukowski, *et al.*, Reference #105.
115. Pandohee J, Holland BJ, Li B, Tsuzuki T, Stevenson PG, Barnett NW, Pearson JR, Jones OAH, Conlan XA. Screening of cannabinoids in industrial-grade hemp using two-dimensional liquid chromatography coupled with acidic potassium permanganate chemiluminescence detection. *Journal of Separation Science* 2015;38(12):2024-2032.
116. Seemann B, Alon T, Tsizin S, Fialkov AB, Amirav A. Electron ionization LC-MS with supersonic molecular beams-the new concept, benefits and applications. *Journal of Mass Spectrometry* 2015;50(11):1252-1263.
117. See Thomas and ElSohly, Reference #1, Figure 4.2, page 69. This figure, which compares the separations of phytocannabinoids on UPLC versus UPC2, is not attributed in the figure caption

or in the associated text; however, it apparently is from the following presentation: Hudalla C. Analytical testing for the cannabis industry: Application of Ultra Performance Convergence Chromatography. Presented at: The 2014 Conference on Small Molecule Science (CoSMoS), Williamsburg, VA:August 11-13, 2014.

118. Aizpurua-Olaizola O, Omar J, Navarro P, Olivares M, Etxebarria N, Usobiaga A. Identification and quantification of cannabinoids in *Cannabis sativa* L. plants by high performance liquid chromatography-mass spectrometry. *Analytical and Bioanalytical Chemistry* 2014;406(29):7549-7560.
119. Jiang G, Stenzel JR, Chen R, Elmashni D. UHPLC/MS analysis of illicit drugs. Chapter 9 in: *Ultra-High Performance Liquid Chromatography and its Applications*, Q. A. Xu, Editor, John Wiley & Sons, Inc., Hoboken, NJ:2013 (pps. 253-269).
120. Mathon C, Duret M, Kohler M, Edder P, Bieri S, Christen P. Multi-targeted screening of botanicals in food supplements by liquid chromatography with tandem mass spectrometry. *Food Chemistry* 2013;138(1):709-717.
121. Wang Q-l, Zhang A-z. UHPLC-MS/MS determination of characteristic cannabinol in hemp food. *Lihua Jianyan, Huaxue Fence* 2013;49(6):720-724.
122. Zhou W-J, Song J-Z, Fu W-W, Tan H-S, Bian Z-X, Xu H-X. Chemical comparison of two dosage forms of hemp seed pills by UHPLC-Q-ToF-MS/MS and multivariate statistical techniques. *Journal of Pharmaceutical and Biomedical Analysis* 2013;84:59-68.
123. Broecker S, Pragst F. Isomerization of cannabidiol and Δ^9 -tetrahydrocannabinol during positive electrospray ionization. In-source hydrogen/deuterium exchange experiments by flow injection hybrid quadrupole-time-of-flight mass spectrometry. *Rapid Communications in Mass Spectrometry* 2012;26(12):1407-1414.
124. Lurie IS, Li L, Toske SG. Hydrophilic interaction chromatography of seized drugs and related compounds with sub-2 μ m particle columns. *Journal of Chromatography A* 2011;1218(52):9336-9344.
125. Stambouli H, El Bouri A, Bouayoun T, El Karni N, Naciri Z, Johar A, Saoura A, Saidi S. Experimentation on industrial hemp crops in Morocco. *Annales de Toxicologie Analytique* 2011;23(1):15-20.
126. Zhang A, Wang Q. Determination of THC in edible vegetable oil by ultra-high performance liquid chromatography-electrospray tandem mass spectrometry. *Shipin Kexue* 2011;32(10):194-198.
127. Jiang G. Use of UHPLC-MS to determine illicit drugs. *American Laboratory* 2010;42(8):40-42.
128. Zhang A, Wang Q, Mo S. Simultaneous determination of Δ^9 -tetrahydrocannabinol, cannabidiol and cannabinol in edible oil using ultra performance liquid chromatography-tandem mass spectrometry. *Sepu* 2010;28(11):1015-1019.
129. Damm M, Rechberger G, Kollroser M, Kappe CO. An evaluation of microwave-assisted derivatization procedures using hyphenated mass spectrometric techniques. *Journal of*

Chromatography A 2009;1216(31):5875-5881.

130. Stenzel JR, Jiang G. Identification of cannabinoids in baked goods by UHPLC/MS. LCGC North America 2009;Suppl:30-31. [Note: Analysis of samples spiked with cannabinoid standards, not with marijuana.]
131. Stolker AAM, Van Schoonhoven J, De Vries AJ, Bobeldijk-Pastorova I, Vaes WHJ, Van den Berg R. Determination of cannabinoids in cannabis products using liquid chromatography-ion trap mass spectrometry. Journal of Chromatography A 2004;1058(1-2):143-151.
132. Cole MD. Analysis of cannabis by supercritical fluid chromatography with ultraviolet detection. Methods In Biotechnology 2000;13:145-148.
133. Radcliffe C, Maguire K, Lockwood B. Applications of supercritical fluid extraction and chromatography in forensic science. Journal of Biochemical and Biophysical Methods 2000;43(1-3):261-272.
134. Ndjoko K, Wolfender JL, Hostettmann K. Analysis of cannabinoids by liquid chromatography - thermospray mass spectrometry and liquid chromatography - tandem mass spectrometry. Chromatographia 1998;47(1/2):72-76.
135. Bäckström B, Cole MD, Carrott MJ, Jones DC, Davidson G, Coleman K. A preliminary study of the analysis of cannabis by supercritical fluid chromatography with atmospheric pressure chemical ionisation mass spectroscopic detection. Science & Justice 1997;37(2):91-97.
136. Lurie IS, Cooper DA, Krull IS. High-performance liquid chromatography using continuous on-line post-elution photoirradiation with subsequent diode-array UV or thermospray mass spectrometry detection. Journal of Chromatography 1993;629(2):143-151.
137. See Thomas and ElSohly, Reference #1.
138. Hillig KW, Mahlberg PG. A chemotaxonomic analysis of cannabinoid variation in Cannabis (*Cannabaceae*). American Journal of Botany 2004;91(6):966-975 (and numerous references cited therein). [Note: Analyses in this study were conducted by GC.]
139. See Happyana, Agnolet, *et al.*, Reference #52.
140. Mechtler K, Bailer B, de Hueber K. Variations of Δ^9 -THC content in single plants of hemp varieties. Industrial Crops and Products 2004;19(1):19-24. [Note: Analyses in this study were conducted by GC/MS.]
141. Csesztregi T, Bovens M, Dujourdy L, Franc A, Nagy J. Sampling of illicit drugs for quantitative analysis - Part III: Sampling plans and sample preparations. Forensic Science International 2014;241:212-229.
142. Bovens M, Csesztregi T, Franc A, Nagy J, Dujourdy L. Sampling of illicit drugs for quantitative analysis - Part II. Study of particle size and its influence on mass reduction. Forensic Science International 2014;234:174-180.

143. Dujourdy L, Csesztregi T, Bovens M, Franc A, Nagy J. Sampling of illicit drugs for quantitative analysis. Part I: Heterogeneity study of illicit drugs in Europe. *Forensic Science International* 2013;231(1-3):249-256. [Note: Does not specifically address cannabis, but provides the background for Parts II and III.]
144. U.S. Food and Drug Administration; Center for Drug Evaluation and Research. Botanical Drug Development - Guidance for Industry (Draft Document for Comment). U.S. FDA; Silver Spring, MD:August, 2015.
145. World Health Organization. Quality Control Methods for Herbal Materials. WHO Press; Geneva, Switzerland:2011.
146. World Health Organization. Quality Control Methods for Medicinal Plant Materials. WHO Press; Geneva, Switzerland:1998.

* * * * *